

Soils of India, Types, Significance

Kuldeep Kumar

Department of Geography Guru Nanak Khalsa College, Karnal

INTRODUCTION

Soils in India include a diverse range of types such as alluvial, black, red, laterite, arid, and forest soils. Check out more Soils of India, Types and Significance.

The soils of India are diverse and shaped by climate, parent material, topography, biological activity, and time. The major soil types include alluvial, black, red, laterite, desert, forest, peaty, and saline soils, each with distinct characteristics and agricultural significance. Alluvial soil is the most fertile and widespread, crucial for agriculture, while black soil is ideal for cotton.

However, soil degradation affects 30% of India's land, caused by unsustainable farming, poor water management, deforestation, and urbanization. Conservation efforts are essential to maintain soil health, ensuring food security and economic stability.

Soils of India

India's diverse geography and climate have resulted in the formation of a wide variety of soils in India, each playing a crucial role in the agricultural landscape. Soil formation is a complex process influenced by several factors, including climate, parent material, topography, biological activity, and time. The type of parent material, such as igneous or sedimentary rocks, determines the soil's composition.

Climate, particularly temperature and rainfall, affect weathering rates and soil fertility, with high rainfall regions experiencing leaching and arid zones leading to less fertile soils. Topography influences soil moisture and vegetation, while biological activity, including the decomposition of organic matter, contributes to soil fertility. Over time, these factors interact, gradually forming distinct soil profiles.

Soils of India Classification

The soils of India are classified based on various criteria, including their physical and chemical properties, as well as their agricultural significance. The Indian Council of Agricultural Research (ICAR) has developed a comprehensive classification system that categorizes Indian soils into eight major groups: alluvial, black, red, laterite, desert (arid), forest, peaty, and saline soils.

Alluvial Soil

Alluvial soil is the most extensive and agriculturally significant soil type in India. Covering about 15 lakh square kilometres or 45.6% of the country's total land area, it forms the foundation of India's agricultural wealth and supports a majority of the population.

Formation and Composition: Alluvial soils are formed by the deposition of sediments (alluvium) by rivers, winds, glaciers, and sea waves. They are rich in humus and nutrients like potash but often lack phosphorus.

Their fertility is further enhanced as they are renewed annually by river floods.

Types of Alluvial Soil: Alluvial soils are categorized into two types:

Old alluvium (Bhangar) is clayey, sticky, and darker with lime nodules.

New alluvium (Khadar) is lighter and deposited in floodplains and deltas.

Distribution: Great Northern Plain (from Punjab to West Bengal and Assam), the lower valleys of the Narmada and Tapi rivers, the east coast deltas, and the northern Gujarat plains through a narrow corridor in Rajasthan.

Characteristics: Alluvial soils range in texture from sandy loam to clay and vary in colour from light grey to ash grey based on maturity.

They are more loamy and clayey in the lower Ganga plain and Brahmaputra valley, with sand content decreasing from west to east.

Crops: Rice, Wheat, Sugarcane, Cotton, Tobacco, Pulses, Oilseeds and Jute.

Black Soil (Regur Soil)

The black soil, also known as Regur or Black Cotton Soil, is primarily found in the Deccan Plateau, covering approximately 5.46 lakh square kilometres (16.6% of India's total geographical area).

Formation and Composition

Black soil, formed from fissure volcanic rock, is ideal for cotton cultivation, earning it the name "black cotton soil." It contains alumina, iron oxide, lime, magnesium carbonates, and poor nitrogen, phosphorus, and organic matter content.

Characteristics: Black soils are highly argillaceous, with over 62% clay, and are excellent at retaining moisture. They have a black colour due to titaniferous magnetite or iron and are ideal for cotton cultivation despite being deficient in essential nutrients.

Crops: Cotton, sugarcane, tobacco, millets (Jowar, sorghum), wheat, oilseeds (groundnut, linseed), pulses (lentils, chickpeas), citrus fruits, sunflowers, turmeric.

Red soils occupy about 3.5 lakh sq km, approximately 10.6% of India's total land area. They are widely distributed across the country, notably in parts of south Bihar, West Bengal, Uttar Pradesh, Aravallis, eastern Rajasthan, and parts of Assam, Nagaland, Manipur, Mizoram, Tripura, and Meghalaya.

Formation and Composition: Red soil forms from the weathering of ancient crystalline rocks, with its red colour due to iron oxide. It is rich in iron but poor in nitrogen, phosphorus, and organic matter content.

Characteristics: Red soils have textures from sandy to loamy, with moderate fertility, suited for dry farming but require irrigation and fertilizers for enhanced productivity.

Crops: Cotton, groundnut, tobacco, and pulses, though less suitable for water-intensive crops due to limited moisture retention.

Laterite Soil

Laterite soils, named after the Latin word "later" meaning brick which hardened upon exposure. Found in regions like South Maharashtra, the Western Ghats in Kerala and Karnataka, parts of Assam, Tamil Nadu, and Western West Bengal (especially in Birbhum district), as well as parts of the Eastern Ghats.

Formation and Composition:

Laterite soils form under high heat and rainfall, causing intense mineral leaching.

They are rich in iron, aluminium, and manganese, giving them reddish to brown hues.

Characteristics: Laterite soils are clayey, compact, and poorly drained, making them less suitable for farming without proper management.

They are poor in essential nutrients like lime, nitrogen, and phosphorus, requiring fertilisers and soil amendments.

Crops: Cashews, tea, coffee, tapioca, rubber, and tropical fruits.

Other Uses: Laterite soils are often used in construction, particularly for making bricks, and roads, and as a building material due to their hardened texture after exposure to air.

Desert (Arid) Soil

Desert soils are primarily found in the arid regions of Rajasthan, Haryana, South Punjab, and parts of Gujarat, as well as near the Indus River and the Aravallis. The Thar Desert is one of the largest stretches of desert soil in India, covering vast areas of northwestern India.

Composition: Desert soils contain high amounts of soluble salts and calcium carbonate, with calcium increasing in deeper layers, making them less fertile.

Distribution: Desert soils cover about 1.42 lakh square kilometres, which is approximately 4.32% of the total land area of India.

These soils are concentrated in the northwestern and western parts of the country.

Characteristics: Saline soils, with low rainfall and poor leaching, are challenging for cultivation without irrigation. Their sandy texture, low organic matter, and poor moisture retention hinder fertility and plant growth.

Crops: Only drought-resistant and salt-tolerant crops like barley, cotton, wheat, millets, and pulses are grown in desert soils.

Forest Soil

Forest soils are predominantly found in the Himalayan region, Western Ghats, and parts of northeastern states like Assam, Nagaland, and Mizoram. These soils are also seen in tropical and subtropical areas, influenced by the type of vegetation and climate.

Composition: Forest soils are rich in organic matter, including humus from decomposed plant material, and are high in essential nutrients like nitrogen, phosphorus, and potassium.

Characteristics: Forest soils vary in texture from sandy loam to clayey, with colours ranging from light to dark brown due to high organic content.

They are fertile but acidic, and deforestation can degrade soil quality, leading to erosion and reduced productivity.

Crops: Tea, coffee, spices, wheat, maize, barley etc.

Peaty Soil

Peaty and marshy soils form in humid regions with excess moisture and waterlogging, slowing plant decomposition. These soils accumulate organic matter under stagnant water and are found in wetland areas like Kottayam and Alappuzha in Kerala, coastal Orissa, Tamil Nadu, Sunderbans in West Bengal, and parts of Bihar and Uttaranchal.

Formation and Composition:

Peaty soils are black due to accumulated decomposed plant material, rich in organic matter and nitrogen, but often lacking in phosphorus and potassium, requiring additional fertilization for crops needing more minerals.

Characteristics: Peaty soils are heavy, spongy, and moisture-retentive, while marshy soils are rich in organic matter and suited for paddy cultivation.

Both are naturally fertile but highly acidic, requiring proper water management for optimal growth.

Crops: Cabbage, broccoli, peas, beans, carrots, radishes, potatoes, lettuce, spinach.

Saline Soil

Saline soils are found in areas like Andhra Pradesh, Karnataka, Bihar, Uttar Pradesh, Haryana, Punjab, Rajasthan, Maharashtra, and parts of Gujarat covering about 68,000 sq km. The accumulation of salts renders soil infertile, with 1.25 million hectares in Uttar Pradesh, 1.21 million hectares in Punjab, and areas around the Gulf of Khambhat in Gujarat affected.

Formation

Saline and alkaline soils are formed in dry regions due to the accumulation of salts like sodium, magnesium, and calcium, primarily through evaporation and weathering of rock and mineral fragments.

Characteristics: Saline and alkaline soils have compact, sandy or silty textures that hinder drainage and root growth. These soils are nutrient-poor, with pale or cracked surfaces, making them infertile for most crops.

It is known by various local names such as reh, kallar, usar, thur, rakar, karl, and chopan.

Crops: Barley, sugar beet, cotton, wheat, rice, sorghum, maize, sunflower, potato, citrus fruits, pomegranate, peas, lentils, cowpea etc.

CONCLUSION

India's diverse topography, from the Himalayan mountains to southern plateaus, fosters a wide range of soils. These varied soils are significant in sustaining diverse agricultural activities, maintaining ecological balance. Diverse soil types support a wide range of agricultural activities, directly sustaining over 40% of India's population. Alluvial soils, in particular, offer highly productive lands essential for farming. Ecosystem Health: Soils are vital for maintaining ecological balance, enabling plant growth that supports animal life and biodiversity. Each soil type fosters specific ecosystems critical for environmental stability. Economic Foundation: Agriculture, dependent on soil fertility, is the backbone of India's economy, supporting 65-70% of the population and influencing food production and economic stability. Carbon Sequestration: Soils are crucial for carbon sequestration, aiding in climate change mitigation. Properly managed soils can capture large amounts of carbon dioxide from the atmosphere, helping to regulate the climate.

REFERENCES

- ABRAHAM J. P. et al, 2013 – A Review of Global Ocean Temperature Observations : Implications for Ocean Heat Content Estimates and Climate Change. Rev. Geophys, 51, 450-483.

- ARMOUR K.C, MARSHALL J. SCOTT J. DONOHOE A and NEWSOM E.R. 2016 – Southern Ocean Warming Delayed by Circumpolar Upwelling and Equatorward Transport, *Nature Geoscience*, 9, 549-554.
- BALMASEDA M. A., TRENBERTH K.E and KALLENE, 2013 – Distinctive Climate Signals in Reanalysis of Global Ocean Heat Content, *Geophys. Res. Lett.* 40, 1754-1759.
- CHEN X. and TUNG K.K., 2014 – Varying Planetary Heat Sink Led to Global-Warming Slowdown and Acceleration, *Science*, 345, 897-903.
- DANABASOGLU G et.al., 2014 – North Atlantic simulations in Coordinated Ocean-ice Reference Experiments phase-II (CORE-II), Part I : Mean states Ocean Modeling, 73, 76-107.
- DECONTO R.M. and POLLARD D. 2016 – Contribution of Antarctica to Past and Future Sea-Level Rise. *Nature*, 531 (7596), pp.591-597.
- DOMINGUES C.M. CHURCH J.A., WHITE N. J., GLECKLER P. J., WIJFFELS S.E., BARKER P. M. AND DUNN J. R. 2008 – Improved Estimates of Upper-Ocean Warming and Multidecadal Sea – Level Rise, *Nature*, 453, 1090 – 1093.
- DRIJFHOUT S.S. BLAKERA, T., JOSEY S. A., NURSER A. J.G. SINHA B. and BALMASEDA M.A., 2014 – Surface Warming Hiatus Caused by Increased Heat Uptake across Multiple Ocean Basins *Geophys Res. Lett.*, 41, 7868-7874.
- DUTEILO, BONING C.W. and OSCHLIES A., 2014 – Variability in Subtropical-Tropical Cells Drives Oxygen Levels in the Tropical Pacific Ocean, *Geophysical Research Letters*, 41 (24) pp. 8926-8934.